

On the completeness of a system of coherent states

A.M. Perelomov

*Institute of Theoretical and Experimental Physics,
117 259 Moscow, USSR*

Abstract

Completeness is proved for some subsystems of a system of coherent states. The linear dependence of states is investigated for the von Neumann type subsystems. A detailed study is made of the case when a regular lattice on the complex α plane with cell area $S = \pi$ corresponds to the states of the system. It is shown that in this case there exists only one linear relationship between the coherent states. This relationship is equivalent to an infinite set of identities. The simplest of these can also be obtained by means of the transformation formulas for θ functions.

1. In the study of a number of problems of quantum mechanics (the theory of measurements [1], quantum optics and radio electronics [2]–[4], and the decay of a quasi-stationary state [5]) it has recently been found useful to employ a system of a so-called coherent states. This is a complete system of states which has found a wide application recently. It is also known that this system of states is supercomplete, i.e., there are the subsystems in this system which are themselves complete. An interesting example of such a system was pointed out by von Neumann [1] in the connection with the problem of the most exact simultaneous measurements of the coordinate and the momentum in quantum mechanics. However, the proof of the completeness of such a subsystem apparently was not published.

The aim of the present note is to prove the completeness of some subsystems of the system of coherent states. In the case of the von Neumann type subsystems, we have also succeeded in investigating of the linear dependence

of the states. It turns out that in the von Neumann case there exists one and only one linear relationship between the states of the subsystems. In the Fock [6] – Bargmann [7] representation, the relationship is equivalent to an infinite set of identities for the functions that are analogous of the θ functions. The simplest of these identities can also be obtained by using of the formulas for the transformation of the θ functions. I did not find the corresponding identities in the mathematical literature in the general case. Finally, the group aspects of this problem are considered briefly.

2. We shall consider an one-dimensional quantum oscillator¹. Let us first recall the well-known properties of such a system (see, for example, [2], [3]). The Hamiltonian of the oscillator has the form²

$$H = \frac{1}{2} (p^2 + x^2) = a^+ a + \frac{1}{2}, \quad (1)$$

where x is the coordinate operators and $p = -i \frac{d}{dx}$ is the momentum operator. The operators

$$a = \frac{1}{\sqrt{2}} (x + ip) = \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right), \quad a^+ = \frac{1}{\sqrt{2}} (x - ip) = \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right)$$

are known as annihilation and creation operators, correspondingly.

The self-adjoint operator H is associated with the system of its eigenvectors $|n\rangle$

$$H |n\rangle = E_n |n\rangle, \quad E_n = n + \frac{1}{2}, \quad n = 0, 1, 2, \dots \quad (2)$$

It is well known that this system is complete and orthogonal.

In many cases, it is more convenient to consider the system of coherent states $|\alpha\rangle$. This is the system of eigenvectors of the annihilation operator a :

$$a |\alpha\rangle = \alpha |\alpha\rangle. \quad (3)$$

Let us mention some of the properties of coherent states (for details, we refer the reader to [2]-[4]). The spectrum of a fills the entire complex plane.

¹ For simplicity, we consider the one-dimensional case. It is readily seen that all results can be directly generalized to the case of N dimensions.

² In this paper we use a system of units in which $\hbar = m = \omega = 1$ ($\hbar$ is the Planck constant, and m and ω are the mass and the frequency of the particle vibrations, correspondingly).

In other words, the state $|\alpha\rangle$ for any complex α can be normalized, i.e., $\langle\alpha|\alpha\rangle = 1$. Expanding such a state with respect to the states $|n\rangle$, we obtain

$$|\alpha\rangle = \exp\left(-\frac{1}{2}|\alpha|^2\right) \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle. \quad (4)$$

The following identity also holds:

$$\frac{1}{\pi} \int d^2\alpha |\alpha\rangle\langle\alpha| = \sum_{n=0}^{\infty} |n\rangle\langle n|, \quad (5)$$

from which it follows that the system of coherent states is complete.

However, the states $|\alpha\rangle$ are not orthogonal to one another. The scalar product of two such states has the form

$$\langle\alpha|\beta\rangle = \exp\left(-\frac{1}{2}(|\alpha|^2 + |\beta|^2 - 2\alpha^*\beta)\right), \quad |\langle\alpha|\beta\rangle|^2 = \exp(-|\alpha - \beta|^2). \quad (6)$$

Equation (5) enables to expand an arbitrary state $|\psi\rangle$ with respect to the states $|\alpha\rangle$:

$$|\psi\rangle = \frac{1}{\pi} \int d^2\alpha \langle\alpha|\psi\rangle |\alpha\rangle. \quad (7)$$

Note that if the coherent state $|\beta\rangle$ is taken as $|\psi\rangle$, then equation (7) defines a linear dependence between the different coherent states. It follows that the system of coherent states is supercomplete, i.e., it contains the subsystems which are complete.

Using (4) we obtain the equation for $\langle\alpha|\psi\rangle$ in (7):

$$\langle\alpha|\psi\rangle = \exp\left(-\frac{1}{2}|\alpha|^2\right) \psi(\alpha^*), \quad (8)$$

where

$$\psi(\alpha) = \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} \langle n|\psi\rangle. \quad (9)$$

At the same time, the inequality $|\langle n|\psi\rangle| \leq 1$ means that the function $\psi(\alpha)$ for the normalization state $|\psi\rangle$ is an entire analytic function of the complex variables α . We also have $|\langle\alpha|\psi\rangle| \leq 1$. Therefore, we obtain a bound on the growth of $\psi(\alpha)$:

$$|\psi(\alpha)| \leq \exp\left(\frac{1}{2}|\alpha|^2\right). \quad (10)$$

The normalization condition can now be written in the form

$$I = \frac{1}{\pi} \int d^2\alpha \exp(-|\alpha|^2) |\psi(\alpha)|^2 = \langle \psi | \psi \rangle. \quad (11)$$

The expansion of an arbitrary state $|\psi\rangle$ with respect to coherent states now takes the form

$$|\psi\rangle = \frac{1}{\pi} \int d^2\alpha \exp\left(-\frac{1}{2}|\alpha|^2\right) \psi(\alpha^*) |\alpha\rangle. \quad (12)$$

Thus, we have established a one-to-one correspondence between the vectors $|\psi\rangle$ of the Hilbert space and the entire analytic functions $\psi(\alpha)$ for which the integral I in (11) is finite. This correspondence is established by equations (9) and (12).

3. Now we can turn to the problem of the completeness of subsystems of coherent states. Following Bargmann [7], we note that since $\psi(\alpha^*)$ is analytic, it is sufficient for its determination to have values of the function $\psi_i = \psi(\alpha_i^*)$ at points α_i^* ($i = 1, 2, 3, \dots$) whose a sequence has a limit point. In particular, if $\psi_i = 0$, (i.e., $\langle \alpha_i | \psi \rangle = 0$), then $\psi(\alpha^*) \equiv 0$ and hence $|\psi\rangle = 0$. This means that the system of states $|\alpha_i\rangle$, $i = 1, 2, 3, \dots$, such that the sequence α_i has a limit point being complete. Note also that these arguments go through for any other system $|\beta_k\rangle$ which differs from the $|\alpha_k\rangle$ system only by the removal of a finite number of states. It follows that any such system is supercomplete.

On the α plane let us now consider a set of points α_i that does not have limit points in a finite part of the plane and the corresponding set of coherent states $\{|\alpha_i\rangle\}$. We can always construct an entire function $\psi(\alpha)$ having zeros at the points α_i^* . At the same time, the state $|\psi\rangle$ constructed by means of equation (12) will be orthogonal to all states $|\alpha_i\rangle$. If, in addition, $|\psi\rangle$ is a square-integrable state, i.e., the integral I in (11) converges, the system $\{|\alpha_i\rangle\}$ is not complete. However, if any state $|\psi\rangle$ that is orthogonal to all the states $|\alpha_i\rangle$ is not square-integrable (i.e., for any entire function with zeros at the points α_i^* the integral I in (11) diverges), the system $\{|\alpha_i\rangle\}$ is complete.

Thus, the question of the completeness of the system $\{|\alpha_i\rangle\}$ is reduced to establishing whether the integral I in (11) converges. In turn, this is determined by the asymptotic behavior of the function $\psi(\alpha)$ as $|\alpha| \rightarrow \infty$. If the order of the growth λ of the entire function $\psi(\alpha)$ is greater than two, the integral diverges; if $\lambda < 2$, the integral converges.

On the other hand, from the theory of entire functions [8], it is well known that the order of growth λ of an entire function $\psi(\alpha)$ with zeros at the points

α_i is not less than the exponent of convergence λ_1 of the sequence α_i , $\lambda \geq \lambda_1$, and it is not difficult to construct functions with $\lambda = \lambda_1$. We recall that the exponent of convergence of a sequence α_i is a number λ_1 such that for the arbitrarily small $\varepsilon > 0$ and $\delta > 0$, the series $\sum_n 1/|\alpha_n|^{\lambda_1+\varepsilon}$ converges and the series $\sum_n 1/|\alpha_n|^{\lambda_1-\delta}$ diverges. At the same time, the series $\sum_n 1/|\alpha_n|^{\lambda_1}$ may either converge or diverge. It follows that for $\lambda_1 > 2$ the system $|\alpha_n\rangle$ is supercomplete and for $\lambda_1 < 2$ such a system is not complete [7].

It remains to consider the more complicated case $\lambda_1 = 2$. If in this case the series $\sum_n 1/|\alpha_n|^2$ converges, then, as follows from the theory of entire functions [8], the function $\psi(\alpha)$ defined by the canonical product has a minimal type. Consequently, the integral I in (11) converges and the system of states $|\alpha_n\rangle$ is not complete.

However, if the series $\sum_n 1/|\alpha_n|^2$ diverges, then to obtain the asymptotic behavior of $\psi(\alpha)$ as $|\alpha| \rightarrow \infty$, we must have a more detailed information about the distribution of the α_n on the α plane. To this end we introduce two quantities Δ and δ . The first of them is the upper density of the distribution of zeros:

$$\Delta = \overline{\lim}_{r \rightarrow \infty} N(r)/r^2, \quad (13)$$

where $N(r)$ is the number of points α_n in a circle of radius r . The quantity δ characterizes the regularity of the distribution of points α_n and is defined as follows:

$$\delta = \overline{\lim}_{r \rightarrow \infty} |\delta(r)|, \quad \delta(r) = \sum_{|\alpha_n| < r} \frac{1}{\alpha_n^2}. \quad (14)$$

Using Lindelöf's theorem [8], we obtain the following result: if $\Delta = 0$ and δ is finite, the system $\{|\alpha_n\rangle\}$ is incomplete; however, if $\Delta = \infty$ or $\delta = \infty$, the system $\{|\alpha_n\rangle\}$ is complete.

The case of $\delta \neq \infty$ and $0 < \Delta < \infty$ requires the more detailed investigation. In this case, $\psi(\alpha)$ is a function of the second growth order ($\lambda = 2$) and a finite type μ . If μ_1 is the minimal type of a function having zeros at the points α_i^* , then for $\mu_1 < 1/2$ the system $\{|\alpha_n\rangle\}$ is incomplete and for $\mu_1 > 1/2$ it is complete. The case $\mu_1 = 1/2$ requires a special investigation.

4. In the general case, we have failed to find an algorithm for the determination of μ_1 in the mathematical literature. However, it is more interesting the case when the points α_i form a regular lattice with a cell area S ($\alpha_{mn} = m\omega_1 + n\omega_2$, where ω_1 and ω_2 are complex numbers with $\text{Im}(\omega_2/\omega_1) \neq 0$, and m and n are integers) can be studied in fair details. Namely, we shall prove the following result.

Assertion 1. *The system of coherent states $\{|\alpha_{mn}\rangle\}$, $\alpha_{mn} = m\omega_1 + n\omega_2$, $\text{Im}(\omega_2/\omega_1) \neq 0$, with a cell area S is not complete if $S > \pi$. It is supercomplete and remains supercomplete if a finite number of states are removed for $S < \pi$. Finally, if $S = \pi$ then the system $\{|\alpha_{mn}\rangle\}$ is complete and it remains complete if a single state is removed but becomes incomplete if any two states are removed.*

Note that for the case of a rectangular lattice, the condition of completeness of the system $\{|\alpha_{mn}\rangle\}$ with $S = \pi$ made an important part in the von Neumann investigation [1] on the question of the most exact simultaneous measurement of the coordinate and momentum. However, evidently the proof of the completeness of such a system was not, apparently, published.

Note also that the α plane is the analog of the phase plane for the classical oscillator, a cell of the phase plane of area $2\pi\hbar$ corresponding to the cell of the α plane of area π . The physical meaning of our assertion becomes clear: the system $\{|\alpha_k\rangle\}$ is complete if we have on the average not less than one coherent state in the cell of the phase plane of area $2\pi\hbar$.

To prove our assertion, we first construct an entire function $\psi(\alpha)$ that has zeros at the lattice sites $\beta_{mn} = \alpha_{mn}^*$. Such a function is well known in the theory of elliptic functions [9], [10] and is usually denoted as $\sigma(\alpha)$:

$$\sigma(\alpha) = \alpha \prod'_{m,n} \left(1 - \frac{\alpha}{\beta_{mn}}\right) \exp\left(\frac{\alpha}{\beta_{mn}} + \frac{1}{2} \frac{\alpha^2}{\beta_{mn}^2}\right). \quad (15)$$

Here the prime indicates that the product in (15) is taken over all points of the lattice with the exception of the point $m = 0, n = 0$. To find the asymptotic behavior of $\sigma(\alpha)$ as $|\alpha| \rightarrow \infty$ we use the identity [9]

$$\sigma(\alpha + \beta_{kl}) = (-1)^{kl+k+l} \exp(\eta_{kl}(\alpha + \beta_{kl}/2)) \sigma(\alpha), \quad (16)$$

where

$$\eta_{kl} = k\eta_1 + l\eta_2, \quad \eta_1 = \zeta(\omega_1^*/2), \quad \eta_2 = \zeta(\omega_2^*/2), \quad (17)$$

and the function $\zeta(\alpha)$ is defined by

$$\zeta(\alpha) = \frac{\sigma'(\alpha)}{\sigma(\alpha)} = \frac{1}{\alpha} + \sum'_{m,n} \left(\frac{1}{\alpha - \beta_{mn}} + \frac{1}{\beta_{mn}} + \frac{\alpha}{\beta_{mn}^2} \right). \quad (18)$$

Hence,

$$|\sigma(\alpha^* + \alpha_{kl}^*)|^2 = \exp\left(\eta_{kl}\alpha^* + \eta_{kl}^*\alpha + \frac{1}{2}(\eta_{kl}\alpha_{kl}^* + \eta_{kl}^*\alpha_{kl})\right) |\sigma(\alpha^*)|^2. \quad (19)$$

Now we turn to a new function $\rho(\alpha, \alpha^*) \geq 0$:

$$|\sigma(\alpha^*)|^2 = \exp\left(\nu\alpha^{*2} + \nu^*\alpha^2 + 2\mu\alpha\alpha^*\right) \rho(\alpha, \alpha^*), \quad (20)$$

and require that this function be doubly periodic, i.e.,

$$\rho(\alpha + \alpha_{m,n}, \alpha^* + \alpha_{mn}^*) = \rho(\alpha, \alpha^*). \quad (21)$$

Hence, we obtain the system of equations for ν and μ :

$$\nu\omega_1^* + \mu\omega_1 = \frac{1}{2}\eta_1, \quad \nu\omega_2^* + \mu\omega_2 = \frac{1}{2}\eta_2, \quad (22)$$

$$\nu\alpha_{mn}^{*2} + \nu^*\alpha_{mn}^2 + 2\mu\alpha_{mn}^*\alpha_{mn} = \frac{1}{2}(\eta_{mn}\alpha_{mn}^* + \eta_{mn}^*\alpha_{mn}). \quad (23)$$

From (22) we find the coefficients ν and μ :

$$\nu = \frac{i}{4S}(\eta_1\omega_2 - \eta_2\omega_1), \quad \mu = \frac{\pi}{2S}. \quad (24)$$

Here $S = \text{Im}(\omega_2\omega_1^*)$ is the cell area.

Note that equation (23) is automatically satisfied. Now it is obvious that the asymptotic behavior of $|\sigma(\alpha^*)|^2$ as $|\alpha| \rightarrow \infty$ is determined by the factor $\exp(\nu\alpha^{*2} + \nu^*\alpha^2 + 2\mu\alpha\alpha^*)$. Thus, the function $\sigma(\alpha^*)$ has the second order and the type $\mu + |\nu|$ (μ and ν are defined in (24)). From equation (20) it is obvious however that $\tilde{\sigma}(\alpha^*) = \sigma(\alpha^*) \exp(-\nu\alpha^{*2})$ is an entire analytic function of the second order and of the minimal possible type μ for the given zeros. Then

$$|\tilde{\sigma}(\alpha^*)|^2 = \rho(\alpha, \alpha^*) \exp(2\nu|\alpha|^2). \quad (25)$$

Using (16) and (24) we obtain a functional equation for $\tilde{\sigma}(\alpha)$:

$$\tilde{\sigma}(\alpha + \alpha_{kl}^*) = (-1)^{kl+k+l} \exp(\mu|\alpha_{kl}|^2) \exp(2\mu\alpha_{kl}\alpha) \tilde{\sigma}(\alpha), \quad \mu = \frac{\pi}{2S}. \quad (26)$$

Invoking arguments similar to those given in the book [10], we conclude that any entire function satisfying this equation is identical with $\tilde{\sigma}(\alpha)$ up to a normalization constant.

The integral I in (11) now takes the form

$$I = \int \rho(\alpha, \alpha^*) \exp\left(\frac{\pi - S}{S}|\alpha|^2\right) d^2\alpha. \quad (27)$$

It is obvious that the integral diverges for $S \leq \pi$ and hence the system $|\alpha_{mn}\rangle$ is complete. Similarly, for $S > \pi$ this system is not complete. Note that if the state $|\alpha_{m_0 n_0}\rangle$ is removed from the system, then we are led to consider the function $(1/(\alpha^* - \alpha_{m_0 n_0}^*)) \tilde{\sigma}(\alpha^*)$. It follows that for $S < \pi$ the system $|\alpha_{mn}\rangle$ remains complete if any finite number of states is removed.

If one of the states $|\alpha_{m_0 n_0}\rangle$ is removed in the case $S = \pi$, the question of the completeness of the system of remaining states reduces to an investigation of the integral

$$I_{m_0 n_0} = \int \frac{\rho(\alpha, \alpha^*)}{|\alpha - \alpha_{m_0 n_0}|^2} d^2 \alpha, \quad (28)$$

which, by virtue of the periodicity of $\rho(\alpha, \alpha^*)$, reduces to I_{00} :

$$I_{m_0 n_0} = I_{00} = \int \frac{d^2 \alpha}{|\alpha|^2} \rho(\alpha, \alpha^*). \quad (29)$$

Here ρ is a non-negative double periodic function all of whose zeros are situated at the points of the lattice $\beta_{mn} = \alpha_{mn}^* = m\omega_1^* + n\omega_2^*$.

Let us estimate I_{00} . Integrating over a region consisting of small nonintersecting circles of radius r_0 with centers γ_{mn} that coincide with the centers of the lattice parallelograms, we obtain

$$I_{00} > \sum'_{m,n} J_{mn}, \quad J_{mn} = \int \rho(\alpha, \alpha^*) \frac{d^2 \alpha}{|\alpha|^2}, \quad |\alpha - \gamma_{mn}| < r_0, \quad (30)$$

but

$$J_{mn} > \frac{\rho_0 \pi r_0^2}{(|\gamma_{mn}| + r_0)^2} > \frac{\pi \rho_0 r_0^2}{4 |\gamma_{mn}|^2},$$

where ρ_0 is the minimum of $\rho(\alpha, \alpha^*)$ in the region $|\alpha - \gamma_{11}| \leq r_0$. Thus,

$$I_{00} > \frac{\pi \rho_0 r_0^2}{4} \sum_{m,n} \frac{1}{|\gamma_{mn}|^2}, \quad \text{i.e.} \quad I_{m_0 n_0} = I_{00} = \infty. \quad (31)$$

It follows that if a single state is removed from the system $|\alpha_{mn}\rangle$, the latter remains complete.

If we remove the states $|\alpha_{m_1 n_1}\rangle$ and $|\alpha_{m_2 n_2}\rangle$, we are led to consider the integral

$$\tilde{I} = \int \frac{\rho(\alpha, \alpha^*) d^2 \alpha}{|\alpha - \alpha_{m_1 n_1}|^2 |\alpha - \alpha_{m_2 n_2}|^2}. \quad (32)$$

Obviously, this integral converges. Hence, the corresponding system now is incomplete. Thus, Assertion 1 is completely proved.

As a complete and minimal system we may take the system of all states $\{|\alpha_{mn}\rangle\}$ with the exception of the state $|0\rangle$. It is well known [11] that in the case of a complete and minimal system, we can always construct an associated complete and minimal system such that these two systems are bi-orthogonal. Suppose that $\{|\tilde{\alpha}_{mn}\rangle\}$ is the system associated with $\{|\alpha_{mn}\rangle\}$ and it is normalized in the usual manner:

$$\langle\alpha_{mn}|\tilde{\alpha}_{kl}\rangle = \delta_{mk}\delta_{nl}. \quad (33)$$

Using (8), we can readily show that the state $|\tilde{\alpha}_{kl}\rangle$ corresponds to the function

$$\tilde{\psi}_{kl}(\alpha^*) = \exp\left(\frac{1}{2}|\alpha_{kl}|^2\right) \frac{\alpha_{kl}^*}{\tilde{\sigma}'(\alpha_{kl}^*)} \frac{\tilde{\sigma}(\alpha^*)}{\alpha^*(\alpha^* - \alpha_{kl}^*)}. \quad (34)$$

The system $|\tilde{\alpha}_{kl}\rangle$ enables to obtain a formal expansion for the arbitrary state $|\psi\rangle$ in a series with respect to the system of states $|\alpha_{mn}\rangle$:

$$|\psi\rangle \sim \sum' c_{mn} |\alpha_{mn}\rangle, \quad (35)$$

where

$$c_{mn} = \langle\tilde{\alpha}_{mn}|\psi\rangle. \quad (36)$$

However, we should remember that we are dealing with a nonorthogonal system and the completeness of the system does not imply, for example, the convergence of the series in (35) [11].³

Taking $|\psi\rangle$ to be the vacuum state $|0\rangle$, we obtain

$$|0\rangle \sim \sum' c_{mn} |\alpha_{mn}\rangle, \quad (37)$$

where

$$c_{mn} = \langle\tilde{\alpha}_{mn}|0\rangle = -\frac{\tilde{\sigma}'^*(0)}{\tilde{\sigma}'^*(\alpha_{mn}^*)} \exp\left(\frac{|\alpha_{mn}|^2}{2}\right) = -(-1)^{m+n+mn}. \quad (38)$$

The relationship (37) can now be rewritten in the form

$$\sum_{m,n} (-1)^{m+n+mn} |\alpha_{mn}\rangle \sim 0. \quad (39)$$

³ Ya.G. Sinai kindly drew my attention to this fact.

To investigate the left-hand side of this relationship we use the Fock [6]–Bargmann [7] representation. We recall that this representation is a concrete realization of the Hilbert space: to the state $|\psi\rangle = \sum c_n |n\rangle$ there corresponds an entire analytic function $\psi(z) = \sum c_n (z^n / \sqrt{n!})$; if $|\psi\rangle = |\alpha\rangle$ is a coherent state, then

$$\psi_\alpha(z) = \exp\left(-\frac{1}{2}|\alpha|^2\right) \exp(\alpha z). \quad (40)$$

In the Fock–Bargmann representation, the relationship (39) takes the form

$$f(z) \equiv \sum_{m,n} (-1)^{m+n+mn} \exp\left(-\frac{1}{2}|\alpha_{mn}|^2 + \alpha_{mn}z\right) \sim 0, \quad (41)$$

where $\alpha_{mn} = m\omega_1 + n\omega_2$, and the cell area is $S = \pi$.

Let us investigate some of the simplest properties of $f(z)$. Obviously, the series (41) converge for all values of z and $f(z)$, and therefore it is an entire function. Further combining in pairs the terms (m, n) and $(-m, -n)$ we see that $f(z)$ is an even function. Let us consider how $f(z)$ varies as a result of a shift of the argument by α_{kl}^* . Under such a shift each term of the series (41) is multiplied by

$$\exp(\alpha_{mn}\alpha_{kl}^*) = \exp\left(\frac{1}{2}(\alpha_{mn}\alpha_{kl}^* + \alpha_{mn}^*\alpha_{kl})\right) \exp\left(\frac{1}{2}(\alpha_{mn}\alpha_{kl}^* - \alpha_{mn}^*\alpha_{kl})\right). \quad (42)$$

However, if the cell area is $S = \pi$, then

$$\exp\left(\frac{1}{2}(\alpha_{mn}\alpha_{kl}^* - \alpha_{mn}^*\alpha_{kl})\right) = (-1)^{ml-nk}, \quad (43)$$

and we obtain

$$\begin{aligned} f(z + \alpha_{kl}^*) &= \sum_{m,n} (-1)^{m+n+mn} (-1)^{ml-nk} \exp\left(-\frac{1}{2}|\alpha_{m-k,n-l}|^2\right) \\ &\times \exp\left(\frac{1}{2}|\alpha_{kl}|^2 + \alpha_{mn}z\right). \end{aligned} \quad (44)$$

Substituting $m \rightarrow m' + k$ and $n \rightarrow n' + l$, we arrive at the identity

$$f(z + \alpha_{kl}^*) = (-1)^{kl+k+l} \exp\left(\frac{1}{2}|\alpha_{kl}|^2\right) \exp(\alpha_{kl}z) f(z). \quad (45)$$

Note that this functional equation is identical with the corresponding equation (26) for the function $\tilde{\sigma}(z)$ when $S = \pi$. Since the solution of this

functional equation is unique (see the remark after equation (26)), we obtain

$$f(z) = C\delta(z), \quad (46)$$

where C is a constant. However, as we have shown above, $f(z)$ is an even function whereas $\tilde{\sigma}(z)$ is an odd function. Therefore, $C = 0$ and consequently $f(z) \equiv 0$.

Thus, we have proved the following assertion.

Assertion 2. *The system of analytic functions $\exp(\alpha_{mn}z)$ ($\alpha_{mn} = m\omega_1 + n\omega_2$, $S = \pi$) is linearly dependent. The equation relating these functions has the form (41).*

Expanding $\exp(\alpha_{mn}z)$ in the Taylor series, we arrive at the identities

$$\sum_{m,n} (-1)^{m+n+mn} \alpha_{mn}^k \exp\left(-\frac{1}{2}|\alpha_{mn}|^2\right) \equiv 0, \quad k = 0, 1, 2, \dots, \quad (47)$$

Let us consider what these identities yield in the simplest cases. If $k = 0$, then (47) can be written in the form

$$\sum_{m,n} (-1)^{m+n+mn} \exp\left(-\frac{\pi}{2}(am^2 + 2bmn + cn^2)\right) \equiv 0, \quad (48)$$

where $ac - b^2 = 1$, $a > 0$, $c > 0$. This identity can be simplified in the case of a rectangular lattice when $b = 0$:

$$\sum_{m,n} (-1)^{m+n+mn} \exp\left(-\frac{\pi}{2}(c^{-1}m^2 + cn^2)\right) = 0. \quad (49)$$

Summing over n and then over m in (49), we obtain

$$\begin{aligned} & \theta_3\left(0, \frac{\tau_1}{2}\right) \theta_3\left(0, \frac{\tau_2}{2}\right) - \theta_3\left(0, \frac{\tau_1}{2}\right) \theta_4\left(0, \frac{\tau_2}{2}\right) \\ & - \theta_4\left(0, \frac{\tau_1}{2}\right) \theta_3\left(0, \frac{\tau_2}{2}\right) - \theta_4\left(0, \frac{\tau_1}{2}\right) \theta_4\left(0, \frac{\tau_2}{2}\right) = 0, \end{aligned} \quad (50)$$

where

$$\tau_1 = ic, \quad \tau_2 = ic^{-1}, \quad c > 0, \quad (51)$$

and the functions $\theta_3(0, \tau)$ and $\theta_4(0, \tau)$ are defined by the equations

$$\theta_3(0, \tau) = \sum_{m=-\infty}^{\infty} \exp(i\pi\tau m^2), \quad \theta_4(0, \tau) = \sum_{m=-\infty}^{\infty} (-1)^m \exp(i\pi\tau m^2). \quad (52)$$

For brevity, we shall write $\theta_i(\tau)$ instead of $\theta_i(0, \tau)$.

We now use the well-known formulas [9], [10]

$$\theta_3(\tau) = \sqrt{\frac{i}{\tau}} \theta_3 \left(-\frac{1}{\tau} \right), \quad \theta_4(\tau) = \sqrt{\frac{i}{\tau}} \theta_2 \left(-\frac{1}{\tau} \right). \quad (53)$$

Equation (50) then becomes

$$\theta_3(2\tau_1) \theta_3(2\tau_2) - \theta_3(2\tau_1) \theta_2(2\tau_2) - \theta_2(2\tau_1) \theta_3(2\tau_2) - \theta_2(2\tau_1) \theta_2(2\tau_2) = 0, \quad (54)$$

where τ_1 and τ_2 are defined in (51).

Using also the doubling formulas [9], [10]

$$\theta_2(2\tau) = \frac{1}{\sqrt{2}} \sqrt{\theta_3^2(\tau) - \theta_4^2(\tau)}, \quad (55)$$

$$\theta_3(2\tau) = \frac{1}{\sqrt{2}} \sqrt{\theta_3^2(\tau) + \theta_4^2(\tau)} \quad (56)$$

and formulas of the type (53)

$$\theta_3(\tau_2) = \sqrt{\frac{i}{\tau_2}} \theta_3 \left(-\frac{1}{\tau_2} \right) = \sqrt{-i\tau_1} \theta_3(\tau_1), \quad (57)$$

$$\theta_4(\tau_2) = \sqrt{\frac{i}{\tau_2}} \theta_2 \left(-\frac{1}{\tau_2} \right) = \sqrt{-i\tau_1} \theta_2(\tau_1),$$

we obtain

$$\begin{aligned} & \sqrt{(\theta_3^2 + \theta_4^2)(\theta_3^2 + \theta_2^2)} - \sqrt{(\theta_3^2 - \theta_4^2)(\theta_3^2 - \theta_2^2)} \\ & - \sqrt{(\theta_3^2 - \theta_4^2)(\theta_3^2 + \theta_2^2)} - \sqrt{(\theta_3^2 + \theta_4^2)(\theta_3^2 - \theta_2^2)} = 0. \end{aligned} \quad (58)$$

Since [9], [10], equation (58) can be transformed into an identity

$$\theta_3^4 = \theta_2^4 + \theta_4^4. \quad (59)$$

Thus, even the proof of the simplest of the identities (47) requires a detailed knowledge of the relationships between θ functions. The proof of the more complicated identities of the type (47) by direct methods is evidently a nontrivial problem.

In this paper, we have considered the case of a single degree of freedom. However, it is readily seen that Eq.(41) can be directly generalized to the case of N degrees of freedom.

5. We are intrigued by the appearance of θ functions in this problem. In this connection, we should like to mention the Cartier paper [12] which establishes a certain connection between the Heisenberg commutation relations for the coordinate and momentum operators and θ functions. Cartier shows that θ functions arise in an investigation of the representation of the group G associated with the commutation relations if one assumes that the representation is induced by a certain representation of a discrete subgroup Γ of G ⁴. However, our approach to the question of θ functions differs from that of Cartier.

Let us briefly consider the relationship between these two approaches. It is readily seen that coherent states can be obtained from the “vacuum state” $|0\rangle$ by means of the so-called displacement operators [2], [3]

$$|\alpha\rangle = D(\alpha) |0\rangle, \quad (60)$$

where the unitary operator $D(\alpha)$ is defined by

$$D(\alpha) = \exp(\alpha a^+ - \alpha^* a), \quad (61)$$

and a^+ and a were defined at the beginning of this paper. The law of multiplication for the operators $D(\alpha)$ has the form

$$D(\alpha_2) D(\alpha_1) = \exp(i \operatorname{Im}(\alpha_2 \alpha_1^*)) D(\alpha_2 + \alpha_1). \quad (62)$$

It follows that the operators $\exp(2\pi i t) D(\alpha)$ form a group G . An element g of this group is defined by a real number t and a complex number α : $g = (t, \alpha)$. The product of two group elements $g = g_2 g_1$ is given by

$$g = (t, \alpha), \quad t = t_2 + t_1 + \frac{1}{2\pi} \operatorname{Im}(\alpha_2 \alpha_1^*), \quad \alpha = \alpha_2 + \alpha_1. \quad (63)$$

Note that the operators $D(\alpha_2)$ and $D(\alpha_1)$ commute if and only if the area of the parallelogram spanned by the vectors α_2 and α_1 is a multiple of π : $S = \operatorname{Im}(\alpha_2 \alpha_1^*) = k\pi$, where k is an integer.

⁴ A number of general questions related to discrete subgroups of continuous groups are discussed in the book by Gel'fand, Graev, and Pyatetskii-Shapiro [13].

Consider the set of operators $D_{np} = D(\alpha_{np})$, where α_{np} is a point of a regular lattice with the cell area $S = \pi$. These operators form a discrete commutative group with multiplication law

$$D_{np}D_{kl} = (-1)^{B(n,p;k,l)} D_{n+k,p+l}, \quad B(n,p;k,l) = nl - pk. \quad (64)$$

Let Γ be the discrete subgroup of G consisting of elements of the form $g = (0, \alpha_{np})$ and $g = (1/2, \alpha_{np})$. The operators $\pm D_{np}$ form a representation of this discrete subgroup and the set of states $|\alpha_{np}\rangle$ forms a basis of a certain representation of the group Γ . Let us try to release the sign factor in (64). To this end, we go over to the new operators

$$D_{np} = (-1)^{F(n,p)} \bar{D}_{np}, \quad (65)$$

and require that

$$\bar{D}_{np} \bar{D}_{kl} = \bar{D}_{n+k,p+l}. \quad (66)$$

This gives the equation for $F(n, p)$:

$$F(n+k, p+l) = F(n, p) + F(k, l) + B(n, p; k, l) \pmod{2} \quad (67)$$

(B is defined in (64)). This equation is identical with equation (71) in [12] for $m = 1$, $N = 1$, and, as it is readily verified, has the solution

$$F(k, l) = kl + k + l. \quad (68)$$

We go over to a new system of states:

$$|\bar{\alpha}_{kl}\rangle = \bar{D}_{kl}|0\rangle = (-1)^{kl+k+l} |\alpha_{kl}\rangle, \quad (69)$$

in which the action of the operator $\bar{D}_{kl}$ has the form

$$\bar{D}_{kl} |\bar{\alpha}_{np}\rangle = |\bar{\alpha}_{k+n, l+p}\rangle. \quad (70)$$

From this equation we can immediately obtain the relationship between the states of the system $|\bar{\alpha}_{np}\rangle$. This is a relationship of the type $\sum c_{np} |\bar{\alpha}_{np}\rangle \sim 0$. As we know, it is unique. Consequently, it can not be changed when the operator $\bar{D}_{kl}$ acts upon it. It is readily seen that the only relationship satisfying this requirement has the form

$$\sum_{m,n} |\bar{\alpha}_{mn}\rangle = \sum_{m,n} (-1)^{mn+m+n} |\alpha_{mn}\rangle \sim 0 \quad (71)$$

which is identical with (39).

Note also that the uniqueness of the solution of the functional equations (26) and (45) for $S = \pi$ follows from the irreducibility of the representation of G induced by the discrete subgroup Γ [12], [13].

I am most indebted to M.I. Graev and Ya.G. Sinai for their interest in the investigation.

References

- [1] von Neumann J., Mathematical Foundations of Quantum Mechanics, Princeton (1955)
- [2] Glauber R.J., Phys. Rev. **130**, 2529; **131**, 2766 (1963)
- [3] Klauder J.R. and Sudarshan E.C., *Fundamentals of Quantum Optics*, Benjamin, New York (1968)
- [4] Zel'dovich Ya.B., Perelomov A.M. and Popov V.S., Zh. Eksp. Teor. Fiz. **55**, 589 (1968); **57**, 196 (1969); Preprints ITEP No. 612, 618 (1968) [in Russian]
- [5] Terent'ev M.V., Preprint ITEP No.773 (1970) [in Russian]
- [6] Fock V.A., Zs. Phys. **49**, 339 (1928)
- [7] Bargmann V., Commun. Pure Appl. Math. **14**, 187 (1961)
- [8] Levin B.Ya., *Distribution of Zeros of Entire Functions*, AMS, Providence, R.I. (1964)
- [9] Erdelyi A. et al (editors), *Higher Transcendental Functions* (California Institute of Technology H. Bateman MS Project), Vol.3, New York (1955)
- [10] Bellman R., *A Brief Introduction to Theta Functions*, Holt, Rinehart and Winston, New York (1961)
- [11] Bari N.K., Uchenye Zapiski MGU **148**, Matem. **4**, 69 (1951)
- [12] Cartier P., Proc. Symp. Pure Math., Vol.9, *Algebraic Groups and Discontinuous Subgroups*, Providence (1966), pp. 361-383

- [13] Gel'fand I.M., Graev M.I. and Pyatetskii-Shapiro I.I., *Theory of Representations and Automorphic Functions*, Nauka (1966) [in Russian]